



# Prediction of Pakistan Super League-2020 Using TOPSIS and Fuzzy TOPSIS Methods

**Muhammad Jafar\*, Faizullah Khan, Amir Naveed**

*Department of Mathematics, Faculty of Basic Sciences Lahore Garrison University, Lahore, Pakistan.*

P A P E R I N F O	A B S T R A C T
<p><b>Chronicle:</b>                      Received: 21 January 2020                      Revised: 05 March 2020                      Accepted: 30 April 2020</p>	<p>We lived in uncertain word and prediction in this uncertain word is a major issue. Prediction in cricket is very complex because there are many factors which are effecting on results. Weather, pitch, Conditions, Home grounds are some of these factors. In this article it is aimed to predict the results of PSL-2020 by using TOPSIS and Fuzzy TOPSIS methods. It will be interesting because first time in PSL history it is going to be held in Pakistan. But we use some serious factor to predict the winner of PSL-2020.</p>
<p><b>Keywords:</b>                      PSL.                      TOPSIS.                      Prediction.                      Cricket.</p>	

## 1. Introduction

Pakistan Super League (PSL) is a professional Twenty20 cricket league in Pakistan contested during February and March every year, with each team play matches in double round robin format. The league is founded by the Pakistan Cricket Board (PCB) in Lahore on 9 September 2015 with initial participation of 5 teams but now there are 6 teams represented by 6 major cities Of Pakistan including Islamabad, Lahore, Multan, Karachi, Peshawar and Quetta.

The commercial rights to the initial franchises were sold for US\$93 million for a ten-years period while the 6th franchise was originally formed in 2017 bought by Schone properties in US\$5.2 million per annum for 8 years contract means the total contract for that team was worth US\$41.6 million that makes it the most expensive team of PSL but unfortunately the contract was terminated by the PCB just after one year due to the non-payment of annual fee by the franchise. Ali Khan Tareen and Taimoor Malik are the new owners of the 6th team as they won the bid for US\$6.2 million per annum for 7 years period.

---

\* Corresponding author  
 E-mail address: naveedjafar@lgu.edu.pk  
 DOI: 10.22105/jfea.2020.249363.1011



**C<sub>3</sub>—match won.** It also an important factor to encourage the teams expectations and moral values. Winning matches shown the previous performance of the team and likely to have to perform well in future on the basis of previous record except of that the pitches, venues and some players are changed for upcoming PSL.

**C<sub>4</sub>—match lost.** It shows the bad performance of the team .The team with most of the lost matches definitely will be considered as the weak team at all but restricted till the expectation as its T20 game and we can't predict with more confidently regarding the strongest and weakest team.

**C<sub>5</sub>—Win % AGE.** Clearly shows the teams combined performance in last 4 editions. As the percentage mostly lies in decimal but in stats it's taken as exact numeric value i.e. instead of taking 58.69%. More the winning percentage shows the team's overall performance in last 4 editions of the PSL.

**C<sub>6</sub>—lost percentage.** Lost percentage shows the weakness of the team. The team with more percentage in matches lost is likely to be consider as the weakest side of the tournament but except of that mostly players are changed or replaced every year so it's not the 100% surety to claim about its performance by just seeing previous record of the weakest team.

**C<sub>7</sub>—all out.** It shows the team's confidence regarding to adopt the pressure in critical situations and also the consistency of the players in difficult situations. More the all outs shows the previous record about their performance in limited overs cricket.

**C<sub>8</sub>—key players.** Key players also plays an important role in match winning .The team with more key players probably have the more chances to win the match or even title. Key players in stats are included with best batting, bowling averages and best economy rates.

**C<sub>9</sub>—best batting averages.** Good batting line up is also a key factor for the team, batting averages taken in stats is >24 means no of best batsmen in a team. More the players with average >24 will increase the chances of winning the match or create a big total on board that helps to win the matches.


**C<sub>9</sub> —bowling strike rate.** Bowling strike rate is the average number of balls bowled per wicket taken, lower the strike rate means more effective a baller is taking wickets quickly. In stats just those players are taken out who have the best strike rates that is consider to be as <19.

**C<sub>11</sub>—economy rate.** Economy rate is the average number of runs conceded for each over bowled. A lower economy rate is preferable. In stats just those bowlers are to be choose out to whom economy rates are less than 8.

*Table 1* shows the statistical behavior of the teams performances. The data is collected up to 4 editions of the PSL.

**Table 1.** Statically behavior of the collected data that is take up to 4 editions of the PSL.

Teams/ Attributes	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
T <sub>1</sub>	31	20	26	16	62	38	6	6	5	2	2
T <sub>2</sub>	31	18	25	18	58	42	5	6	6	4	4
T <sub>3</sub>	27	8	17	23	43	57	4	6	6	3	3
T <sub>4</sub>	36	15	27	19	59	41	4	4	6	2	2
T <sub>5</sub>	12	5	7	12	37	63	6	4	4	1	4
T <sub>6</sub>	24	5	10	24	31	69	7	4	4	2	3




**Fig. 1.** Graphical representation of the above mentioned data.

### 3. Calculations

TOPSIS technique is used to calculate an MCDM problem. So the weights are assigned according to the importance of each of the attribute.

<b>Weight</b>	0.15	0.08	0.12	0.03	0.09	0.08	0.02	0.17	0.16	0.08	0.02
<b>Attributes</b>	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>



**Fig. 2.** Graphical behavior of the above mentioned weights to the attributes.

There is a great demand of game prediction for that many of the mathematical formulas are to be used for prediction by using MCDM techniques that is used to choose the best one amongst multiple criteria's.

### 3.1. TOPSIS Prediction

TOPSIS, Technique for order of preference by similarity to ideal solution, a tool for calculating the best one from the collected data. TOPSIS technique is applied to calculate the weights by given data that is collected up to 4 editions of the PSL which is easily available on "www.cricinfo.com" that is considered to be one of the best cricketing website in the world. To apply the technique some important attributes that are considered to play an important role in match or title winning including Fifties, chased, matches won, matches lost, win percentage, lost percentage, best batting average, best bowling strike rate, best economy rate with 6 teams T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub>, and T<sub>6</sub> as decision makers are taken.

**Step 1.** Calculate normalized matrix.

$$X_{ij} = \frac{X_{ij}}{\sqrt{\sum_{j=1}^n X^2_{ij}}}$$

<b>Weight</b>	<b>0.15</b>	<b>0.08</b>	<b>0.12</b>	<b>0.03</b>	<b>0.09</b>
<b>T/C</b>	<b>C<sub>1</sub></b>	<b>C<sub>2</sub></b>	<b>C<sub>3</sub></b>	<b>C<sub>4</sub></b>	<b>C<sub>5</sub></b>
<b>T1</b>	0.453777	0.613428	0.523360293	0.341899075	0.508470092
<b>T2</b>	0.453777	0.552085	0.503231051	0.384636459	0.47566557
<b>T3</b>	0.395225	0.245371	0.342197115	0.49147992	0.352648612
<b>T4</b>	0.526967	0.460071	0.543489536	0.406005151	0.483866701
<b>T5</b>	0.175656	0.153357	0.140904694	0.256424306	0.303441829
<b>T6</b>	0.351311	0.153357	0.201292421	0.512848612	0.254235046
<b>0.08</b>	<b>0.02</b>	<b>0.17</b>	<b>0.16</b>	<b>0.08</b>	<b>0.02</b>
<b>C<sub>6</sub></b>	<b>C<sub>7</sub></b>	<b>C<sub>8</sub></b>	<b>C<sub>9</sub></b>	<b>C<sub>10</sub></b>	<b>C<sub>11</sub></b>
0.292584827	0.449719013	0.480384461	0.389249472	0.324442842	0.262612866
0.32338323	0.374765844	0.480384461	0.467099366	0.648885685	0.525225731
0.43887724	0.299812676	0.480384461	0.467099366	0.486664263	0.393919299
0.315683629	0.299812676	0.320256308	0.467099366	0.324442842	0.262612866
0.485074845	0.449719013	0.320256308	0.311399578	0.162221421	0.525225731
0.531272449	0.524672182	0.320256308	0.311399578	0.324442842	0.393919299

**Step 2.** Calculate the weighted normalized matrix.

$$V_{ij} = X_{ij} \times W_j$$

Weight	0.15	0.08	0.12	0.03	0.09
T/C	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>
<b>T1</b>	0.068067	0.049074	0.062803235	0.010256972	0.045762308
<b>T2</b>	0.068067	0.044167	0.060387726	0.011539094	0.042809901
<b>T3</b>	0.059284	0.01963	0.041063654	0.014744398	0.031738375
<b>T4</b>	0.079045	0.036806	0.065218744	0.012180155	0.043548003
<b>T5</b>	0.026348	0.012269	0.016908563	0.007692729	0.027309765
<b>T6</b>	0.052697	0.012269	0.02415509	0.015385458	0.022881154
<b>0.08</b>	<b>0.02</b>	<b>0.17</b>	<b>0.16</b>	<b>0.08</b>	<b>0.02</b>
C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
0.023406786	0.00899438	0.081665358	0.062279916	0.025955427	0.005252257
0.025870658	0.007495317	0.081665358	0.074735899	0.051910855	0.010504515
0.035110179	0.005996254	0.081665358	0.074735899	0.038933141	0.007878386
0.02525469	0.005996254	0.054443572	0.074735899	0.025955427	0.005252257
0.038805988	0.00899438	0.054443572	0.049823932	0.012977714	0.010504515
0.042501796	0.010493444	0.054443572	0.049823932	0.025955427	0.007878386

**Step 3.** Calculate the ideal best and ideal worst values.

	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
<b>V<sup>+</sup></b>	0.079	0.049	0.0652	0.0076	0.0457	0.0234	0.0059	0.0816	0.0747	0.0519	0.0105
<b>+</b>	045	074	18744	92729	62308	06786	96254	65358	35899	10855	04515
<b>V<sup>-</sup></b>	0.026	0.012	0.0169	0.0153	0.0228	0.0425	0.0104	0.0544	0.0498	0.0129	0.0052
<b>-</b>	348	269	08563	85458	81154	01796	93444	43572	23932	77714	52257

**Step 4.** Calculate the Euclidean distance from the idea best.

$$Si^+ = [(\sum_{j=1}^m V_{ij} - V_j^+)^2]^{0.5}$$

**Step 5.** Calculate the Euclidean distance from the ideal worst.

$$Si^- = [(\sum_{j=1}^m V_{ij} - V_j^-)^2]^{0.5}$$

**Step 6.** Calculate the performance score.

$$P_i = \frac{Si^-}{Si^+ + Si^-}$$

Teams	Si+	Si-	Pi	Rank
<b>Quetta Gladiators</b>	0.032	0.085	0.728429	2
<b>Islamabad United</b>	0.014	0.091	0.865348	1
<b>Karachi Kings</b>	0.049	0.063	0.560988	4
<b>Peshawar Zalmi</b>	0.04	0.085	0.679043	3
<b>Multan Sultan</b>	0.1	0.011	0.099888	6
<b>Lahore Qalandars</b>	0.082	0.03	0.269933	5




Fig. 3. Graphical representation of the final results against each team.

### 3.2. Calculations via Fuzzy TOPSIS

**Step 1.** Assign the fuzzy numbers to the group of decision makers on the basis of attributes regarding each of the criteria.

T/C	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
T <sub>1</sub>	8,6,7	9,7,8	8,6,7	8,6,7	9,7,8	2,3,5	6,5,4	9,7,8	8,6,7	8,6,7	8,6,7
T <sub>2</sub>	8,6,7	8,6,7	6,5,4	6,5,4	8,6,7	6,5,4	2,3,5	6,5,4	6,5,4	6,5,4	8,6,7
T <sub>3</sub>	6,5,4	2,3,5	6,5,4	2,3,5	2,3,5	6,5,4	2,3,5	9,7,8	8,6,7	6,5,4	6,5,4
T <sub>4</sub>	9,7,8	6,5,4	9,7,8	6,5,4	8,6,7	2,3,5	2,3,5	6,5,4	8,6,7	2,3,5	2,3,5
T <sub>5</sub>	1,1,2	1,1,2	2,3,5	2,3,5	2,3,5	8,6,7	8,6,7	6,5,4	2,3,5	1,1,2	8,6,7
T <sub>6</sub>	2,3,5	1,1,2	1,1,2	1,1,2	1,1,2	9,7,8	9,7,8	6,5,4	2,3,5	2,3,5	6,5,4

#### Decision Maker 2.

	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
T <sub>1</sub>	9,7,8	9,7,8	8,6,7	8,6,7	9,7,8	1,1,2	6,5,4	9,7,8	9,7,8	9,7,8	8,6,7
T <sub>2</sub>	8,6,7	8,6,7	6,5,4	6,5,4	8,6,7	2,3,5	1,1,2	6,5,4	8,6,7	8,6,7	6,5,4
T <sub>3</sub>	6,5,4	1,1,2	6,5,4	2,3,5	2,3,5	6,5,4	2,3,5	9,7,8	8,6,7	8,6,7	6,5,4
T <sub>4</sub>	8,6,7	6,5,4	8,6,7	6,5,4	8,6,7	2,3,5	2,3,5	6,5,4	6,5,4	2,3,5	2,3,5
T <sub>5</sub>	1,1,2	1,1,2	2,3,5	6,5,4	2,3,5	8,6,7	8,6,7	6,5,4	2,3,5	1,1,2	8,6,7
T <sub>6</sub>	2,3,5	1,1,2	1,1,2	6,5,4	1,1,2	8,6,7	9,7,8	6,5,4	2,3,5	2,3,5	8,6,7

#### Decision Maker 3.

	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
T <sub>1</sub>	8,6,7	8,6,7	8,6,7	9,7,8	8,6,7	1,1,2	2,3,5	9,7,8	9,7,8	8,6,7	6,5,4
T <sub>2</sub>	8,6,7	8,6,7	6,5,4	8,6,7	8,6,7	2,3,5	1,1,2	8,6,7	6,5,4	8,6,7	8,6,7
T <sub>3</sub>	6,5,4	2,3,5	6,5,4	2,3,5	2,3,5	6,5,4	2,3,5	9,7,8	9,7,8	6,5,4	6,5,4
T <sub>4</sub>	9,7,8	6,5,4	9,7,8	6,5,4	8,6,7	2,3,5	2,3,5	6,5,4	8,6,7	2,3,5	2,3,5
T <sub>5</sub>	2,3,5	6,5,4	2,3,5	2,3,5	2,3,5	8,6,7	8,6,7	6,5,4	2,3,5	1,1,2	8,6,7
T <sub>6</sub>	1,1,2	2,3,5	1,1,2	9,7,8	1,1,2	9,7,8	9,7,8	6,5,4	2,3,5	2,3,5	6,5,4

**Step 2.** Construct the combined decision matrix by using the following formula.

$$\tilde{x}_{ij} = (a_{ij}, b_{ij}, c_{ij}) \text{ where } a_{ij} = \min_k \{a_{ij}^k\}, b_{ij} = \frac{1}{k} \sum_{k=1}^k b_{ij}^k, c_{ij} = \max_k \{c_{ij}^k\}.$$

	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
T <sub>1</sub>	8,6.33,8	8,6.66,8	8,3,7	8,6.33,8	8,6.66,8	1,1.66,5	2,4.33,5	9,7,8	8,6.66,8	8,6.33,8	6,5.66,7
T <sub>2</sub>	8,6,7	8,6,7	6,5,4	6,5.33,7	8,6,7	2,3.66,5	1,1.66,5	6,5.33,7	6,5.33,7	6,5.66,7	6,5.66,7
T <sub>3</sub>	6,3,4	1,2.33,5	6,5,4	2,3,5	2,3,5	6,5,4	2,3,5	9,7,8	8,6.33,7	6,5.33,7	6,5,4
T <sub>4</sub>	8,6.66,8	6,5,4	8,6.66,8	6,5,4	8,6,7	2,3,5	2,3,5	6,5,4	6,5.66,7	2,3,5	2,3,5
T <sub>5</sub>	1,1.66,5	1,2.33,4	2,3,5	2,3.66,5	2,3,5	8,6,7	8,6,7	6,5,4	2,3,5	1,1,2	8,6,7
T <sub>6</sub>	1,2.33,5	1,1.66,5	1,1,2	1,4.33,8	1,1,2	8,7,8	9,7,8	6,5,4	2,3,5	2,3,5	6,5.33,7

**Step 3.** Compute the normalized fuzzy decision matrix by using formulae.

$$r_{ij} = \left(\frac{a_{ij}}{c_j^+}, \frac{b_{ij}}{c_j^+}, \frac{c_{ij}}{c_j^+}\right) \text{ and } c_j^+ = \max_i \{c_{ij}\} \text{ (Beneficial criteria).}$$

$$r_{ij} = \left(\frac{a_j^-}{c_{ij}^-}, \frac{a_j^-}{b_{ij}^-}, \frac{a_j^-}{a_{ij}^-}\right) \text{ and } a_j^- = \min_i \{a_{ij}\} \text{ (Non beneficial criteria).}$$

	Bnf	Bnf	Bnf	Non Bnf	Bnf
Wght	8,6,9	6,5,4	7,5,8	6,5,4	6,3,5
T/C	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>
T <sub>1</sub>	1,0.79125,1	1,0.8325,1	1,0.375,0.875	0.125,0.1579,0.125	1,0.8325,1
T <sub>2</sub>	1,0.75,1	1,0.75,0.875	0.75,0.625,0.5	0.1428,0.1876,0.166	1,0.75,0.875
T <sub>3</sub>	0.75,0.375,0.5	0.125,0.291,0.625	0.75,0.625,0.5	0.2,0.33,0.5	0.25,0.375,0.625
T <sub>4</sub>	1,0.8325,1	0.75,0.626,0.5	1,0.8325,1	0.25,0.2,0.166	1,0.75,0.875
T <sub>5</sub>	0.125,0.2075,0.625	0.125,0.291,0.5	0.25,0.375,0.625	0.2,0.27,0.166	0.25,0.375,0.625
T <sub>6</sub>	0.125,0.291,0.625	0.125,0.2075,0.625	0.125,0.125,0.25	0.125,0.230,1	0.125,0.125,0.25
Non Bnf	Non Bnf	Bnf	Bnf	Bnf	Bnf
3,4,7	4,5,6	9,7,9	9,6,8	6,5,7	7,6,8
C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
0.2,0.602,1	0.2,0.230,0.5	1.125,0.875,1	1,0.8325,1	1,0.791,1	0.8571,0.8085,1
0.2,0.2732,0.5	0.2,0.6024,1	0.75,0.666,0.875	0.75,0.666,0.875	0.75,0.7075,0.875	0.8571,0.8085,1
0.25,0.22,0.166	0.2,0.33,0.5	1.125,0.875,1	0.75,0.666,0.875	0.75,0.666,0.875	0.8571,0.7142,0.5715
0.2,0.33,0.5	0.2,0.33,0.5	0.75,0.625,0.5	0.75,0.7075,0.875	0.25,0.375,0.625	0.285,0.4285,0.7142
0.1425,0.166,0.125	0.1425,0.166,0.125	0.75,0.625,0.5	0.25,0.375,0.625	0.125,0.125,0.25	1.1428,0.8571,1
0.125,0.1425,0.125	0.12,0.14,0.111	0.75,0.625,0.5	0.25,0.375,0.625	0.25,0.375,0.625	0.8571,0.7614,1

**Step 4.** Calculate weighted normalized fuzzy decision matrix.

$$v_{ij} = r_{ij} \times w_j.$$

Such that  $A_1 \otimes A_2 = (a_1, b_1, c_1) \otimes (a_2, b_2, c_2) = (a_1 \times a_2, b_1 \times b_2, c_1 \times c_2).$



	<b>Bnf</b>	<b>Bnf</b>	<b>Bnf</b>	<b>Non Bnf</b>	<b>Bnf</b>
<b>Wght</b>	<b>8,6,9</b>	<b>6,5,4</b>	<b>7,5,8</b>	<b>6,5,4</b>	<b>6,3,5</b>
<b>T/C</b>	<b>C<sub>1</sub></b>	<b>C<sub>2</sub></b>	<b>C<sub>3</sub></b>	<b>C<sub>4</sub></b>	<b>C<sub>5</sub></b>
<b>T1</b>	8,4.7475,9	6,4.1625,4	7,1.87,6.7	0.75,0.7895,0.5	6,2.49755,5
<b>T2</b>	8,4,5,9	6,3.75,3.5	5.25,3.125,4	0.8568,0.938,0.664	6,2.25,4.375
<b>T3</b>	6,2.25,4.5	0.75,1.4575,2.5	5.25,3.125,4	1.2,1.65,2	1.5,1.125,3.1285
<b>T4</b>	8,4.995,9	4.5,3.125,2	7,4.1625,8	1.5,1,0.64	6,2.25,4.125
<b>T5</b>	1,1.245,5.626	0.75,1.4550,2	1.75,1.875,5	1.2,1.35,2	1.5,1.125,3.125
<b>T6</b>	1,1.7475,5.625	0.75,1.037,2.5	0.875,0.625,2	0.75,1.15,4	0.75,0.375,1.25
<b>Non Bnf</b>	<b>Non Bnf</b>	<b>Bnf</b>	<b>Bnf</b>	<b>Bnf</b>	<b>Bnf</b>
<b>3,4,7</b>	<b>4,5,6</b>	<b>9,7,9</b>	<b>9,6,8</b>	<b>6,5,7</b>	<b>7,6,8</b>
<b>C<sub>6</sub></b>	<b>C<sub>7</sub></b>	<b>C<sub>8</sub></b>	<b>C<sub>9</sub></b>	<b>C<sub>10</sub></b>	<b>C<sub>11</sub></b>
0.6,2.408,7	0.8,1.15,3	10.125,6.125,9	9,4.995,8	6,3.955,7	5.99,4.851,8
0.6,1.092,3.5	0.8,3.012,6	6.75,4.662,7.875	6.75,3.996,7	4.5,3.5375,6.125	5.99,4.851,8
0.75,0.88,1.162	0.8,1.65,3	9,5.39,7.92	9,4.74,7	4.5,3.3,6.125	5.25,3.75,4
0.6,1.32,3.5	0.8,1.65,3	5.94,3.85,3.96	5.58,4.62,7	1.5,1.875,4.37	1.75,2.25,5
0.42,0.64,0.84	0.56,0.80,0.72	5.94,3.85,3.96	2.25,2.25,5	0.75,0.625,1.75	7,4,5,7
0.36,0.57,0.84	0.48,0.70,0.66	5.94,3.85,3.96	2.25,2.25,5	1.5,1.875,4.375	5.25,3.96,7

**Step 5.** Compute the Fuzzy Positive Ideal Solution (FPIS) and Fuzzy Negative Ideal Solution (FNIS) by given formula.

$$A^+ = (v_1^+, v_2^+ \dots v_n^+) \text{ where } v_j^+ = \max_i \{v_{ij}\}.$$

$$A^- = (v_1^-, v_2^- \dots v_n^-) \text{ where } v_j^- = \min_i \{v_{ij}\}.$$

	<b>C<sub>1</sub></b>	<b>C<sub>2</sub></b>	<b>C<sub>3</sub></b>	<b>C<sub>4</sub></b>	<b>C<sub>5</sub></b>	<b>C<sub>6</sub></b>	<b>C<sub>7</sub></b>	<b>C<sub>8</sub></b>	<b>C<sub>9</sub></b>	<b>C<sub>10</sub></b>	<b>C<sub>11</sub></b>
<b>FPIS</b>	8,4.99	6,4.16	7,4.16	0.75,	6,2.497	0.6,2.4	0.8,3.0	10.125,	9,4.9	6,3.955	5.99,4
<b>= A<sup>+</sup></b>	5,9	25,4	25,8	1.15,	55,5	08,7	12,6	6.125,9	95,8	,7	.851,8
<b>FNIS</b>	1,1.24	0.75,1.	0.875,	0.75,	0.75,0.	0.36,0.	0.48,0.	5.94,3.	2.25,	0.75,0.	1.75,2
<b>= A<sup>-</sup></b>	5,5.62	037,2.	0.625,	1.15,	375,1.2	57,0.8	70,0.6	85,3.96	2.25,	625,1.7	.25,5
	6	5	2	4	5	4	6		5	5	

**Step 6.** Calculate the distance from each alternative to the FPIS and then FNIS by using distance formula as mentioned below.

$$d(x,y) = \sqrt{\frac{1}{3} [(a_1 - a_2)^2 + (b_1 - b_2)^2 + (c_1 - c_2)^2]}.$$

### Distance from FPIS.

T/C	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>
<b>T1</b>	0.142894192	0	1.521573336	2.0314166	0
<b>T2</b>	0.285788383	0.37423533	2.5909397	1.967769485	0.388107051
<b>T3</b>	3.25499744	3.830831348	2.5909397	1.2232095	2.916048
<b>T4</b>	0	1.562733	0	1.9887454	0.505181
<b>T5</b>	4.981511685	3.711749374	3.73252176	1.1913192	2.924003035
<b>T6</b>	4.862676946	3.632320207	5.355045323	0	3.912826
C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
0	2.03854883	0	0	0	0
2.158846606	0	2.220857117	1.53411223	1.031169441	0
3.485184261	1.902195574	0.995364255	0.595825758	1.071548723	2.433091381
2.116108378	1.902195574	4.003808603	2.068576403	3.24002572	3.353744226
3.70152329	3.308017735	4.003808603	4.549634601	4.698010217	0.84524178
3.714002513	3.364686414	4.003808603	4.549634601	3.24002572	0.883455526

### Distance from FNIS.

T/C	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>
<b>T1</b>	4.921090064	3.63232	4.514999077	2.0314166	3.92131859
<b>T2</b>	4.864116946	3.70463	3.12999	1.967769485	3.664666
<b>T3</b>	3.015399	0.241188	3.12999	1.2232095	1.245493
<b>T4</b>	4.98048	2.49481	5.355045323	1.9887454	3.6214
<b>T5</b>	0	0.376364	1.942133	1.1913192	1.243734
<b>T6</b>	0.29011	0	0	0	0
C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>
3.724002513	1.388104223	4.003808603	4.549634601	4.698010217	3.353744226
2.424574464	3.364686414	2.355325101	3.016538634	3.727634695	3.494042
0.34248309	1.46975	3.023066875	4.311326169	3.667821788	2.273030283
1.60163582	1.46975	0	2.627153085	1.731030522	0
0.053229	0.081649658	0	0	0	3.494042549
0	0	0	0	1.733553052	2.52811524

**Step 7.** Calculate the  $d_i^+$  and  $d_i^-$  by using the formula mentioned below.

$$d_i^+ = \sum_{j=1}^n d(v_{ij}, v_j^+) \quad d_i^- = \sum_{j=1}^n d(v_{ij}, v_j^-).$$

**Step 8.** Calculate the Closeness Coefficient (CC<sub>i</sub>) for each alternative.

$$CC_i = d_i^- / (d_i^+ + d_i^-).$$

Teams	di+	di-	Cci	RANK
<b>Quetta Gladiators</b>	5.734432958	40.73844871	0.8766069	1
<b>Islamabad United</b>	12.55182534	35.71397374	0.739943695	2
<b>Karachi Kings</b>	24.29923594	23.9427577	0.496305312	4
<b>Peshawar Zalmi</b>	20.7411183	25.87005015	0.555018272	3
<b>Multan Sultan</b>	37.64734128	8.382471407	0.182109614	5
<b>Lahore Qalandars</b>	37.51848185	4.551778292	0.108194679	6

So T<sub>1</sub> (Quetta Gladiator) has the most chances to win PSL-2020.




Fig. 4. Graphical representation of the final results by FTOPSIS.

#### 4. Conclusion

The purpose of this research is to predict about the expected winner of PSL-5-2020. Although it's really a problematic task to predict about the game like this especially when we talk about limited overs cricket. Many of the factors including pitch, weather conditions, availability of the key players directly impacts on the game, except of all that this research paper is based on the previous records of all the teams as well as availability of the key players for the upcoming edition of the PSL. So by collected records and mathematically point of view the results shows that team Islamabad United probably have the more chances to win the PSL-2020 by TOPSIS technique but fuzzy TOPSIS shows slightly different results with winning chances of another strongest and well deserved winning team Quetta Gladiators as well as the teams Multan Sultans and Lahore Qalandars have the lowest chances to win the title.

#### References

- [1] Ahmed, F., Deb, K., & Jindal, A. (2013). Multi-objective optimization and decision making approaches to cricket team selection. *Applied soft computing*, 13(1), 402-414.
- [2] Bhattacharya, R., Gill, P. S., & Swartz, T. B. (2011). Duckworth–Lewis and twenty20 cricket. *Journal of the operational research society*, 62(11), 1951-1957.
- [3] Davis, J., Perera, H., & Swartz, T. B. (2015). Player evaluation in Twenty20 cricket. *Journal of sports analytics*, 1(1), 19-31.
- [4] Perera, H. P., & Swartz, T. B. (2013). Resource estimation in T20 cricket. *IMA journal of management mathematics*, 24(3), 337-347.