

Paper Type: Research Paper



A Study on Fundamentals of Refined Intuitionistic Fuzzy Set with Some Properties

Atiqe Ur Rahman^{1,*} , Muhammad Rayees Ahmad¹, Muhammad Saeed¹, Muhammad Ahsan¹,
Muhammad Arshad¹, Muhammad Ihsan¹

Department of Mathematics, University of Management and Technology, Lahore, Pakistan; aurkhhb@gmail.com;
rayeesmalik.ravian@gmail.com; muhammad.saeed@umt.edu.pk; ahsan1826@gmail.com; maakhhb84@gmail.com; mihkhhb@gmail.com.

Citation:



Montazeri, F. Z. (2020). An overview of data envelopment analysis models in fuzzy stochastic environments. *Journal of fuzzy extension and application*, 1 (4), 272-292.

Received: 01/08/2020

Reviewed: 11/09/2020

Revised: 04/11/2020


Accept: 03/12/2020

Abstract

Zadeh conceptualized the theory of fuzzy set to provide a tool for the basis of the theory of possibility. Atanassov extended this theory with the introduction of intuitionistic fuzzy set. Smarandache introduced the concept of refined intuitionistic fuzzy set by further subdivision of membership and non-membership value. The meagerness regarding the allocation of a single membership and non-membership value to any object under consideration is addressed with this novel refinement. In this study, this novel idea is utilized to characterize the essential elements e.g. subset, equal set, null set, and complement set, for refined intuitionistic fuzzy set. Moreover, their basic set theoretic operations like union, intersection, extended intersection, restricted union, restricted intersection, and restricted difference, are conceptualized. Furthermore, some basic laws are also discussed with the help of an illustrative example in each case for vivid understanding.

Keywords: Fuzzy set, Intuitionistic fuzzy set, Refined intuitionistic fuzzy set.

1 | Introduction

 Licensee **Journal of Fuzzy Extension and Applications**. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

Zadeh [1] introduced the concept of fuzzy set for the first time in 1965 which covers all weak aspects of the classical set theory. In fuzzy set, the membership value is allocated from the interval $[0, 1]$ to all the elements of the universe under consideration. Zadeh [2] used his own concept as a basis for a theory of possibility. Dubois and Prade [3] and [4] established relationship between fuzzy sets and probability theories and also derive monotonicity property for algebraic operations performed between random set-valued variables. Ranking fuzzy numbers in the setting of possibility theory was done by Dubois and Prade [5]. This concept was used by Liang et al. in data analysis, similarities measures in fuzzy sets were discussed by Beg and Ashraf [6]-[8].



Corresponding Author: aurkhhb@gmail.com



10.22105/JFEA.2021.281500.1061

Set difference and symmetric difference of fuzzy sets were established by Vemuri et al., after that, Neog and Sut [9] extended the work to complement of an extended fuzzy set. A lot of work is done by researchers in fuzzy mathematics and its hybrids [10]-[16].

In some real life situations, the values are in the form of intervals due to which it is hard to allocate a membership value to the element of the universe of discourse. Therefore, the concept of interval-valued is introduced which proved a very powerful tool in this area.

In 1986, Atanassov [17] and [18] introduced the concept of intuitionistic fuzzy set in which the membership value and non-membership value is allocated from the interval $[0,1]$ to all the elements of the universe under consideration. It is the generalization of the fuzzy set. The invention of intuitionistic fuzzy set proved very important tool for researchers. Ejegwa et al. [19] discussed about operations, algebra, model operators and normalization on intuitionistic fuzzy sets. Szmidt and Kacprzyk [20] gave geometrical representation of an intuitionistic fuzzy set is a point of departure for our proposal of distances between intuitionistic fuzzy sets and also discussed properties. Szmidt and Kacprzyk [21] also discussed about non-probabilistic-type entropy measure for these sets and used it for geometrical interpretation of intuitionistic fuzzy sets. Proposed measure in terms of the ratio of intuitionistic fuzzy cardinalities was also defined and discussed. Ersoy and Davvaz [22] discussed the basic definitions and properties of intuitionistic fuzzy Γ -hyperideals of a Γ -semi-hyperring with examples are introduced and described some characterizations of Artinian and Noetherian Γ -semi hyper ring. Bustince and Burillo [23] proved that vague sets are intuitionistic fuzzy sets. A lot of work is done by researchers in intuitionistic fuzzy environment and its hybrids [24]-[33].

In 2019, Smarandache defined the concept of refined intuitionistic fuzzy set [34]. In this paper, we extend the concept to refined intuitionistic fuzzy set and defined some fundamental concepts and aggregation operations of refined intuitionistic fuzzy set.

Imprecision is a critical viewpoint in any decision making procedure. Various tools have been invented to deal with the uncertain environment. Perhaps the important tool in managing with imprecision is intuitionistic fuzzy sets. Besides, the most significant thing is that in real life scenario, it is not sufficient to allocate a single membership and non-membership value to any object under consideration. This inadequacy is addressed with the introduction of refined intuitionistic fuzzy set. Having motivation from this novel concept, essential elements, set theoretic operations and basic laws are characterized for refined intuitionistic fuzzy set in this work.

The remaining article is outlined in such a way that the Section 2 recalls some basic definitions along with illustrative example. Section 3 explains basic notions of Refined Intuitionistic Fuzzy Set (RIFS) including subset, equal set, null set and complement set along with their examples for the clear understanding. Section 4 explains the aggregation operations of RIFS with the help of example, Section 5 gives some basic laws of RIFS and in the last, Section 6 concludes the work and gives the future directions.

2 | Preliminaries

In this section, some basic concepts of Fuzzy Set (FS), Intuitionistic Fuzzy Set (IFS) and RIFS are discussed.

Let us consider \tilde{U} be a universal set, N be a set of natural numbers, I represent the interval $[0,1]$, T_{η}^{ω} denotes the degree of sub-truth of type $\omega = 1, 2, 3, \dots, \alpha$ and F_{η}^{λ} denotes the degree of sub-falsity of type $\lambda = 1, 2, 3, \dots, \beta$ such that α and β are natural numbers. An illustrative example is considered to understand these entire basic concepts throughout the paper.

Definition 1. [1, 2] The fuzzy set $\tilde{\eta}_f = \{ \langle \delta, \alpha_{\tilde{\eta}_f}(\delta) \rangle \mid \delta \in \tilde{U} \}$ on \tilde{U} such that $\alpha_{\tilde{\eta}_f}(\delta): \tilde{U} \rightarrow \tilde{I}$ where $\alpha_{\tilde{\eta}_f}(\delta)$ describes the membership of $\delta \in \tilde{U}$.




Fig. 1. Representation of fuzzy set.

Example 1. Hamna wants to purchase a dress for farewell party event of her university. She expected to purchase such dress which meets her desired requirements according to the event. Let $\tilde{U} = \{\tilde{B}_1, \tilde{B}_2, \tilde{B}_3, \tilde{B}_4\}$, be different well-known brands of clothes in Pakistan such that

\tilde{B}_1 = Ideas Gul Ahmad;
 \tilde{B}_2 = Khaadi;
 \tilde{B}_3 = Nishat Linen;
 \tilde{B}_4 = Junaid jamshaid.

Then fuzzy set $\tilde{\eta}_f$ on the universe \tilde{U} is written in such a way that $\tilde{\eta}_f = \left\{ \langle \tilde{B}_1, 0.45 \rangle, \langle \tilde{B}_2, 0.57 \rangle, \langle \tilde{B}_3, 0.6 \rangle, \langle \tilde{B}_4, 0.64 \rangle \right\}$.

Definition 2. [18]. An IFS $\tilde{\eta}_{IFS}$ on \tilde{U} is given by $\tilde{\eta}_{IFS} = \{ \langle \delta, T_{\tilde{\eta}}(\delta), F_{\tilde{\eta}}(\delta) \rangle \mid \delta \in \tilde{U} \}$,

where $T_{\tilde{\eta}}(\delta), F_{\tilde{\eta}}(\delta): \tilde{U} \rightarrow P([0,1])$, respectively, with the condition $\sup T_{\tilde{\eta}}(\delta) + \sup F_{\tilde{\eta}}(\delta) \leq 1$.

Example 2. Consider the illustrative example, and then the intuitionistic fuzzy set $\tilde{\eta}_{IFS}$ on the universe \tilde{U} is given as $\tilde{\eta}_{IFS} = \{ \langle \tilde{B}_1, 0.75, 0.14 \rangle, \langle \tilde{B}_2, 0.57, 0.2 \rangle, \langle \tilde{B}_3, 0.6, 0.3 \rangle, \langle \tilde{B}_4, 0.64, 0.16 \rangle \}$.

Definition 3. [34] A RIFS $\tilde{\eta}_{RIFS}$ on \tilde{U} is given by $\tilde{\eta}_{RIFS} = \{ \langle \delta, T_{\tilde{\eta}}^{\omega}(\delta), F_{\tilde{\eta}}^{\lambda}(\delta) \rangle : \omega \in N_1^{\alpha}, \lambda \in N_1^{\beta}, \alpha + \beta \geq 3, \delta \in \tilde{U} \}$, where $\alpha, \beta \in \tilde{I}$ such that $T_{\tilde{\eta}}^{\omega}, F_{\tilde{\eta}}^{\lambda} \subseteq \tilde{I}$, respectively, with the condition

$$\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}}^{\omega}(\delta) + \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}}^{\lambda}(\delta) \leq 1.$$

It is denoted by $(\tilde{\delta}, \tilde{G})$, where $\tilde{G} = (T_{\tilde{\eta}}^{\omega}, F_{\tilde{\eta}}^{\lambda})$.

Example 3. Consider the illustrative example, then the RIFS $\tilde{\eta}_{RIFS}$ can be written in such a way that

$$\tilde{\eta}_{RIFS} = \{ \langle \tilde{B}_1, (0.5, 0.4), (0.3, 0.25) \rangle, \langle \tilde{B}_2, (0.35, 0.3), (0.15, 0.1) \rangle, \langle \tilde{B}_3, (0.35, 0.25), (0.3, 0.2) \rangle, \langle \tilde{B}_4, (0.6, 0.1), (0.12, 0.2) \rangle \}.$$

3| Basic Notions of RIFS

In this section, some basic notions of subset, equal sets, null set and complement set for RIFS are introduced.

Definition 4. Refined intuitionistic fuzzy subset

Let $\check{\eta}_{1RIFS} = (\check{\delta}, \check{G}_1)$ and $\check{\eta}_{2RIFS} = (\check{\delta}, \check{G}_2)$ be two RIFS, then $\check{\eta}_{1RIFS} \subseteq \check{\eta}_{2RIFS}$, if

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}) \leq \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}), \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}) \geq \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \quad \forall \quad \check{\delta} \in \check{U}.$$

Remark 1. If

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}) < \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}), \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}) > \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \quad \forall \quad \check{\delta} \in \check{U}.$$

Then it is denoted by $(\check{\delta}, \check{G}_1) \subset (\check{\delta}, \check{G}_2)$.

Suppose $(\check{\delta}, \check{G}_1^i) \subset (\check{\delta}, \check{G}_2^i)$ be two families of RIFS, then $(\check{\delta}, \check{G}_1^i)$ is called family of refined intuitionistic fuzzy subset of $(\check{\delta}, \check{G}_2^i)$, if $\check{G}_1^i \subset \check{G}_2^i$ and

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}) < \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}), \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}) > \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}), \quad \forall \quad \check{\delta} \in \check{U}.$$

We denote it by $(\check{\delta}, \check{G}_1^i) \subset (\check{\delta}, \check{G}_2^i) \quad \forall \quad i = 1, 2, 3, \dots, n$.

Example 4. Consider the illustrative example, let $\check{\eta}_{1RIFS}$ and $\check{\eta}_{2RIFS}$ be two RIFS such that

$$\check{\eta}_{1RIFS} = \{ \langle \check{B}_1, (0.35, 0.1), (0.22, 0.19) \rangle, \langle \check{B}_2, (0.25, 0.03), (0.15, 0.19) \rangle, \langle \check{B}_3, (0.2, 0.1), (0.2, 0.24) \rangle, \langle \check{B}_4, (0.3, 0.4), (0.06, 0.04) \rangle \},$$

and

$$\check{\eta}_{2RIFS} = \{ \langle \check{B}_1, (0.38, 0.11), (0.2, 0.14) \rangle, \langle \check{B}_2, (0.45, 0.04), (0.1, 0.14) \rangle, \langle \check{B}_3, (0.3, 0.2), (0.01, 0.06) \rangle, \langle \check{B}_4, (0.31, 0.41), (0.01, 0.011) \rangle \}.$$

Then from above equations, it is clear that $\check{\eta}_{1RIFS} \subseteq \check{\eta}_{2RIFS}$.

Definition 5. Equal refined intuitionistic fuzzy sets

Let $\check{\eta}_{1RIFS} = (\check{\delta}, \check{G}_1)$ and $\check{\eta}_{2RIFS} = (\check{\delta}, \check{G}_2)$ be two RIFS, then $\check{\eta}_{1RIFS} = \check{\eta}_{2RIFS}$ if $\check{\eta}_{1RIFS} \subseteq \check{\eta}_{2RIFS}$ and $\check{\eta}_{2RIFS} \subseteq \check{\eta}_{1RIFS}$.

Example 5. Consider the illustrative example, let $\check{\eta}_{1RIFS}$ and $\check{\eta}_{2RIFS}$ be two RIFS such that

$$\check{\eta}_{1RIFS} = (\check{\delta}, \check{G}_1) = \{ \langle \check{B}_1, (0.4, 0.5), (0.03, 0.04) \rangle, \langle \check{B}_2, (0.5, 0.4), (0.05, 0.04) \rangle, \langle \check{B}_3, (0.5, 0.2), (0.01, 0.06) \rangle, \langle \check{B}_4, (0.3, 0.4), (0.06, 0.04) \rangle \},$$

and

$$\check{\eta}_{2_{RIFS}} = (\check{\delta}, \check{\zeta}_2) = \{ \langle \check{B}_1, (0.4, 0.5), (0.03, 0.04) \rangle, \langle \check{B}_2, (0.5, 0.4), (0.05, 0.04) \rangle, \langle \check{B}_3, (0.5, 0.2), (0.01, 0.06) \rangle, \langle \check{B}_4, (0.3, 0.4), (0.06, 0.04) \rangle \}.$$

Then from above equations, it is clear that $\check{\eta}_{1_{RIFS}} = \check{\eta}_{2_{RIFS}}$.

Definition 6. Null refined intuitionistic fuzzy set

Let RIFS $(\check{\delta}, \check{\zeta})$ is said to be null RIFS if

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}}^{\omega}(\check{\delta}) = 0, \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}}^{\lambda}(\check{\delta}) = 0, \quad \forall \check{\delta} \in \check{\mathcal{U}}.$$

It is denoted by $(\check{\delta}, \check{\zeta})_{null}$.

Example 6. Consider the illustrative example, the null RIFS is given as

$$(\check{\delta}, \check{\zeta}) = \{ \langle \check{B}_1, (0, 0), (0, 0) \rangle, \langle \check{B}_2, (0, 0), (0, 0) \rangle, \langle \check{B}_3, (0, 0), (0, 0) \rangle, \langle \check{B}_4, (0, 0), (0, 0) \rangle \}.$$

Definition 7. Complement of refined intuitionistic fuzzy set

The complement of RIFS $(\check{\delta}, \check{\zeta})$ is denoted by $(\check{\delta}, \check{\zeta}^c)$ and is defined that if

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}^c}^{\omega}(\check{\delta}) = \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}}^{\lambda}(\check{\delta}), \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}^c}^{\lambda}(\check{\delta}) = \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}}^{\omega}(\check{\delta}), \quad \forall \check{\delta} \in \check{\mathcal{U}}.$$

Remark 2. The complement of family of RIFS $(\check{\delta}, \check{\zeta}^c)$ is denoted by $(\check{\delta}, \check{\zeta}^c)$ and is defined in a way that if

$$\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_i^c}^{\omega}(\check{\delta}) = \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}}^{\lambda}(\check{\delta}), \quad \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_i^c}^{\lambda}(\check{\delta}) = \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}}^{\omega}(\check{\delta}), \quad \forall i = 1, 2, 3, \dots, n.$$

Example 7. Consider the illustrative example, if there is a RIFS $\check{\eta}_{RIFS}$ given as

$$\check{\eta}_{RIFS} = \{ \langle \check{B}_1, (0.2, 0.1), (0.3, 0.35) \rangle, \langle \check{B}_2, (0.05, 0.34), (0.45, 0.04) \rangle, \langle \check{B}_3, (0.01, 0.6), (0.1, 0.02) \rangle, \langle \check{B}_4, (0.3, 0.04), (0.12, 0.2) \rangle \}.$$

Then the complement of RIFS $\check{\eta}_{RIFS}$ given as

$$\check{\eta}_{RIFS}^c = \{ \langle \check{B}_1, (0.3, 0.35), (0.2, 0.1) \rangle, \langle \check{B}_2, (0.45, 0.04), (0.05, 0.34) \rangle, \langle \check{B}_3, (0.1, 0.02), (0.01, 0.6) \rangle, \langle \check{B}_4, (0.12, 0.2), (0.3, 0.04) \rangle \}.$$

4 | Aggregation Operators of RIFS

In this section, union, intersection, extended intersection, restricted union, restricted intersection and restricted difference of RIFS is defined with the help of illustrative example.

Definition 8. Union of two RIFS

The union of two RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is denoted by $(\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2)$ and it is defined as $(\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) = (\check{\delta}, \check{Y})$, where $\check{Y} = \check{G}_1 \cup \check{G}_2$, and truth and false membership of $(\check{\delta}, \check{Y})$ is defined in such a way that

$$T_{\check{Y}}(\check{\delta}) = \max \left(\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}), \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}) \right),$$

$$F_{\check{Y}}(\check{\delta}) = \min \left(\sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}), \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \right).$$

Remark 3. The union of two families of RIFS $(\check{\delta}, \check{G}_1^i)$ and $(\check{\delta}, \check{G}_2^i)$ is denoted by $(\check{\delta}, \check{G}_1^i) \cup (\check{\delta}, \check{G}_2^i)$ and it is defined as $(\check{\delta}, \check{G}_1^i) \cup (\check{\delta}, \check{G}_2^i) = (\check{\delta}, \check{Y}^i)$, where $\check{Y}^i = \check{G}_1^i \cup \check{G}_2^i$, $i = 1, 2, 3, \dots, n$, and truth and false membership of $(\check{\delta}, \check{Y}^i)$ is defined in such a way that

$$T_{\check{Y}^i}(\check{\delta}) = \max \left(\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}), \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}) \right),$$

$$F_{\check{Y}^i}(\check{\delta}) = \min \left(\sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}), \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \right).$$

Example 8. Consider the illustrative example, suppose that

$$(\check{\delta}, \check{G}_1) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \},$$

and

$$(\check{\delta}, \check{G}_2) = \{ \langle \check{B}_1, (0.2, 0.3), (0.3, 0.15) \rangle, \langle \check{B}_2, (0.32, 0.38), (0.1, 0.04) \rangle, \langle \check{B}_3, (0.01, 0.16), (0.5, 0.2) \rangle, \langle \check{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \},$$

be two RIFS. Then the union of RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is given as

$$(\check{\delta}, \check{Y}) = \{ \langle \check{B}_1, (0.2, 0.3), (0.24, 0.1) \rangle, \langle \check{B}_2, (0.32, 0.38), (0.1, 0.04) \rangle, \langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \}.$$

Definition 9. Intersection of two RIFS

The intersection of two RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is denoted by $(\check{\delta}, \check{G}_1) \cap (\check{\delta}, \check{G}_2)$ and it is defined as $(\check{\delta}, \check{G}_1) \cap (\check{\delta}, \check{G}_2) = (\check{\delta}, \check{Y})$, where $\check{Y} = \check{G}_1 \cap \check{G}_2$, and truth and false membership of $(\check{\delta}, \check{Y})$ is defined in such a way that

$$T_{\tilde{Y}}(\delta) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_1}^{\omega}(\delta), \sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_2}^{\omega}(\delta) \right),$$

$$F_{\tilde{Y}}(\delta) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_1}^{\lambda}(\delta), \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_2}^{\lambda}(\delta) \right).$$

Remark 4. The intersection of two families of RIFS (δ, \tilde{G}_1^i) and (δ, \tilde{G}_2^i) is denoted by $(\delta, \tilde{G}_1^i) \cap (\delta, \tilde{G}_2^i)$ and it is defined as $(\delta, \tilde{G}_1^i) \cap (\delta, \tilde{G}_2^i) = (\delta, \tilde{Y}^i)$, where $\tilde{Y}^i = \tilde{G}_1^i \cap \tilde{G}_2^i$, $i = 1, 2, 3, \dots, n$, and truth and false membership of (δ, \tilde{Y}^i) is defined in such a way that

$$T_{\tilde{Y}^i}(\delta) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_1}^{\omega}(\delta), \sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_2}^{\omega}(\delta) \right),$$

$$F_{\tilde{Y}^i}(\delta) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_1}^{\lambda}(\delta), \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_2}^{\lambda}(\delta) \right).$$

Example 9. Consider the illustrative example, suppose that

$$\begin{aligned} (\delta, \tilde{G}_1) = \{ < \tilde{B}_1, (0.13, 0.19), (0.24, 0.1) >, \\ < \tilde{B}_2, (0.2, 0.25), (0.15, 0.24) >, \\ < \tilde{B}_3, (0.1, 0.36), (0.34, 0.12) >, < \tilde{B}_4, (0.16, 0.14), (0.23, 0.37) > \}, \end{aligned}$$

and

$$\begin{aligned} (\delta, \tilde{G}_2) = \{ < \tilde{B}_1, (0.2, 0.3), (0.3, 0.15) >, \\ < \tilde{B}_2, (0.32, 0.38), (0.1, 0.04) >, \\ < \tilde{B}_3, (0.01, 0.16), (0.5, 0.2) >, < \tilde{B}_4, (0.26, 0.15), (0.12, 0.2) > \}, \end{aligned}$$

be two RIFS. Then the intersection of RIFS (δ, \tilde{G}_1) and (δ, \tilde{G}_2) is given as

$$\begin{aligned} (\delta, \tilde{Y}) = \{ < \tilde{B}_1, (0.13, 0.19), (0.24, 0.1) >, \\ < \tilde{B}_2, (0.2, 0.25), (0.15, 0.24) >, \\ < \tilde{B}_3, (0.01, 0.16), (0.5, 0.2) >, < \tilde{B}_4, (0.16, 0.14), (0.23, 0.37) > \}. \end{aligned}$$

Definition 10. Extended intersection of two RIFS

The intersection of two RIFS (δ, \tilde{G}_1) and (δ, \tilde{G}_2) is denoted by $(\delta, \tilde{G}_1) \cap_{\epsilon} (\delta, \tilde{G}_2)$ and it is defined as $(\delta, \tilde{G}_1) \cap_{\epsilon} (\delta, \tilde{G}_2) = (\delta, \tilde{Y})$, where $\tilde{Y} = \tilde{G}_1 \cup \tilde{G}_2$, and truth and false membership of (δ, \tilde{Y}) is defined in such a way that

$$T_{\tilde{Y}}(\delta) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_1}^{\omega}(\delta), \sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_2}^{\omega}(\delta) \right),$$

$$F_{\tilde{Y}}(\delta) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_1}^{\lambda}(\delta), \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_2}^{\lambda}(\delta) \right).$$

Remark 5. The extended intersection of two families of RIFS (δ, \tilde{G}_1^i) and (δ, \tilde{G}_2^i) is denoted by $(\delta, \tilde{G}_1^i \cap_{\epsilon} \tilde{G}_2^i)$ and it is defined as $(\delta, \tilde{G}_1^i) \cap_{\epsilon} (\delta, \tilde{G}_2^i) = (\delta, \tilde{Y}^i)$, where $\tilde{Y}^i = \tilde{G}_1^i \cap \tilde{G}_2^i$, $i = 1, 2, 3, \dots, n$, and truth and false membership of (δ, \tilde{Y}^i) is defined in such a way that

$$T_{\tilde{Y}^i}(\delta) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_1}^{\omega}(\delta), \sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_2}^{\omega}(\delta) \right),$$

$$F_{\tilde{Y}^i}(\delta) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_1}^{\lambda}(\delta), \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_2}^{\lambda}(\delta) \right).$$

Example 10. Consider the illustrative example, suppose that

$$\begin{aligned} (\delta, \tilde{G}_1) &= \{ \langle \tilde{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \\ &\langle \tilde{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \\ &\langle \tilde{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle \}, \end{aligned}$$

and

$(\delta, \tilde{G}_2) = \{ \langle \tilde{B}_3, (0.01, 0.16), (0.5, 0.2) \rangle, \langle \tilde{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \}$, be two RIFS. Then the extended intersection of RIFS (δ, \tilde{G}_1) and (δ, \tilde{G}_2) is given as

$$\begin{aligned} (\delta, \tilde{Y}) &= \{ \langle \tilde{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \langle \tilde{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \\ &\langle \tilde{B}_3, (0.01, 0.16), (0.5, 0.2) \rangle, \langle \tilde{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \}. \end{aligned}$$

Definition 11. Restricted union of two RIFS

The restricted union of two RIFS (δ, \tilde{G}_1) and (δ, \tilde{G}_2) is denoted by $(\delta, \tilde{G}_1) \cup_R (\delta, \tilde{G}_2)$ and it is defined as $(\delta, \tilde{G}_1) \cup_R (\delta, \tilde{G}_2) = (\delta, \tilde{Y})$, where $\tilde{Y} = \tilde{G}_1 \cap_R \tilde{G}_2$, and truth and false membership of (δ, \tilde{Y}) is defined in such a way that

$$T_{\tilde{Y}}(\delta) = \max \left(\sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_1}^{\omega}(\delta), \sum_{\omega=1}^{\alpha} \sup T_{\tilde{\eta}_2}^{\omega}(\delta) \right).$$

$$F_{\tilde{Y}}(\delta) = \min \left(\sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_1}^{\lambda}(\delta), \sum_{\lambda=1}^{\beta} \sup F_{\tilde{\eta}_2}^{\lambda}(\delta) \right).$$

Remark 6. The restricted union of two families of RIFS (δ, \tilde{G}_1^i) and (δ, \tilde{G}_2^i) is denoted by

$(\delta, \tilde{G}_1^i) \cup_R (\delta, \tilde{G}_2^i)$ and it is defined as $(\delta, \tilde{G}_1^i) \cup_R (\delta, \tilde{G}_2^i) = (\delta, \tilde{Y}^i)$, where $\tilde{Y}^i = \tilde{G}_1^i \cap_R \tilde{G}_2^i$, $i = 1, 2, 3, \dots, n$, and truth and false membership of (δ, \tilde{Y}^i) is defined in such a way that

$$T_{\check{\Upsilon}i}(\check{\delta}) = \max \left(\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}), \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}) \right),$$

$$F_{\check{\Upsilon}i}(\check{\delta}) = \min \left(\sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}), \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \right).$$

Example 11. Consider the illustrative example, suppose that

$$(\check{\delta}, \check{G}_1) = \{ < \check{B}_1, (0.13, 0.19), (0.24, 0.1) >,$$

$$< \check{B}_2, (0.2, 0.25), (0.15, 0.24) >,$$

$$< \check{B}_3, (0.1, 0.36), (0.34, 0.12) > \},$$

and $(\check{\delta}, \check{G}_2) = \{ < \check{B}_3, (0.01, 0.16), (0.5, 0.2) >, < \check{B}_4, (0.26, 0.15), (0.12, 0.2) > \}$, be two RIFS. Then the restricted union of RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is given as

$$(\check{\delta}, \check{\Upsilon}) = \{ < \check{B}_3, (0.1, 0.36), (0.34, 0.12) > \}.$$

Definition 12. Restricted intersection of two RIFS

The restricted intersection of two RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is denoted by $(\check{\delta}, \check{G}_1) \cap_R (\check{\delta}, \check{G}_2)$ and it is defined as $(\check{\delta}, \check{G}_1) \cap_R (\check{\delta}, \check{G}_2) = (\check{\delta}, \check{\Upsilon})$, where $\check{\Upsilon} = \check{G}_1 \cap_R \check{G}_2$, and truth and false membership of $(\check{\delta}, \check{\Upsilon})$ is defined in such a way that

$$T_{\check{\Upsilon}}(\check{\delta}) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}), \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}) \right),$$

$$F_{\check{\Upsilon}}(\check{\delta}) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}), \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \right).$$

Remark 7. The restricted intersection of two families of RIFS $(\check{\delta}, \check{G}_1^i)$ and $(\check{\delta}, \check{G}_2^i)$ is denoted by

$(\check{\delta}, \check{G}_1^i) \cap_R (\check{\delta}, \check{G}_2^i)$ and it is defined as $(\check{\delta}, \check{G}_1^i) \cap_R (\check{\delta}, \check{G}_2^i) = (\check{\delta}, \check{\Upsilon}^i)$, where $\check{\Upsilon}^i = \check{G}_1^i \cap_R \check{G}_2^i$, $i = 1, 2, 3, \dots, n$, and truth and false membership of $(\check{\delta}, \check{\Upsilon}^i)$ is defined in such a way that

$$T_{\check{\Upsilon}i}(\check{\delta}) = \min \left(\sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_1}^{\omega}(\check{\delta}), \sum_{\omega=1}^{\alpha} \sup T_{\check{\eta}_2}^{\omega}(\check{\delta}) \right),$$

$$F_{\check{\Upsilon}i}(\check{\delta}) = \max \left(\sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_1}^{\lambda}(\check{\delta}), \sum_{\lambda=1}^{\beta} \sup F_{\check{\eta}_2}^{\lambda}(\check{\delta}) \right).$$

Example 12. Consider the illustrative example, suppose that

$$(\check{\delta}, \check{G}_1) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \\ \langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \\ \langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle \},$$

$$\text{and } (\check{\delta}, \check{G}_2) = \{ \langle \check{B}_2, (0.32, 0.38), (0.1, 0.04) \rangle, \langle \check{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \},$$

be two RIFS. Then the restricted intersection of RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is given as

$$(\check{\delta}, \check{Y}) = \{ \langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle \}.$$

Definition 13. Restricted difference of two RIFS

The restricted difference of two RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is denoted by $(\check{\delta}, \check{G}_1)_{-R}(\check{\delta}, \check{G}_2)$ and it is defined as $(\check{\delta}, \check{G}_1)_{-R}(\check{\delta}, \check{G}_2) = (\check{\delta}, \check{Y})$, where $\check{Y} = \check{G}_1 -_R \check{G}_2$.

Example 13. Consider the illustrative example, suppose that

$$(\check{\delta}, \check{G}_1) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \\ \langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \\ \langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \},$$

and

$$(\check{\delta}, \check{G}_2) = \{ \langle \check{B}_1, (0.2, 0.3), (0.3, 0.15) \rangle, \langle \check{B}_2, (0.32, 0.38), (0.1, 0.04) \rangle, \\ \langle \check{B}_3, (0.01, 0.16), (0.5, 0.2) \rangle \},$$

be two RIFS. Then the restricted difference of RIFS $(\check{\delta}, \check{G}_1)$ and $(\check{\delta}, \check{G}_2)$ is given as

$$(\check{\delta}, \check{Y}) = \{ \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \}.$$

5 | Some Basic Laws of RIFS

In this section, we prove some basic fundamental laws including idempotent law, identity law, domination law, De-Morgan law and commutative law with the help of illustrative example.

5.1 | Idempotent Law

$$(\check{\delta}, \check{G}) \cup (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cup_R (\check{\delta}, \check{G}).$$

$$(\check{\delta}, \check{G}) \cap (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cap_\epsilon (\check{\delta}, \check{G}).$$

Example 14. To prove (1) law, we consider illustrative example. For this, suppose that

$$(\check{\delta}, \check{G}) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle,$$

$$\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle,$$

$$\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \}.$$

One can observe

$$(\check{\delta}, \check{G}) \cup (\check{\delta}, \check{G}) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle,$$

$$\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle,$$

$$\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \} = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cup_R (\check{\delta}, \check{G}).$$

Similarly, we can prove (2).

5.2 | Identity Law

$$(\check{\delta}, \check{G}) \cup \emptyset = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cup_R \emptyset.$$

$$(\check{\delta}, \check{G}) \cap \check{U} = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cap_{\epsilon} \check{U}.$$

Example 15. To prove (1) law, we consider illustrative example. For this, suppose that

$$(\check{\delta}, \check{G}) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle,$$

$$\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle,$$

$$\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \}.$$

One can observe

$$(\check{\delta}, \check{G}) \cup \emptyset = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle,$$

$$\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle,$$

$$\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \} = (\check{\delta}, \check{G}) = (\check{\delta}, \check{G}) \cup_R \emptyset.$$

Similarly, we can *Prove* (2).

5.3 | Domination Law

$$(\check{\delta}, \check{G}) \cup \check{U} = \check{U} = (\check{\delta}, \check{G}) \cup_R \check{U}.$$

$$(\check{\delta}, \check{G}) \cap \emptyset = \emptyset = (\check{\delta}, \check{G}) \cap_{\epsilon} \emptyset.$$

Example 16. To *Prove* (1) law, we consider illustrative example. For this, suppose that

$$(\check{\delta}, \check{G}) = \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle,$$

$$\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle,$$

$$\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \}.$$

One can observe

$$\begin{aligned}
 (\check{\delta}, \check{G}) \cup \check{U} &= \{ \langle \check{B}_1, (0.13, 0.19), (0.24, 0.1) \rangle, \\
 &\langle \check{B}_2, (0.2, 0.25), (0.15, 0.24) \rangle, \\
 &\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.16, 0.14), (0.23, 0.37) \rangle \} \cup \check{U} \\
 &= \check{U} = (\check{\delta}, \check{G}) \cup_R \check{U}.
 \end{aligned}$$

Similarly, we can *Prove* (2).

5.4 | De-Morgan Law

$$\begin{aligned}
 \left((\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) \right)^c &= (\check{\delta}, \check{G}_1)^c \cap_\epsilon (\check{\delta}, \check{G}_2)^c. \\
 \left((\check{\delta}, \check{G}_1) \cap_\epsilon (\check{\delta}, \check{G}_2) \right)^c &= (\check{\delta}, \check{G}_1)^c \cup (\check{\delta}, \check{G}_2)^c.
 \end{aligned}$$

Example 17. To prove (1) law, we consider illustrative example. For this, suppose that L.H.S is

$$\begin{aligned}
 (\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) &= \{ \langle \check{B}_1, (0.2, 0.3), (0.24, 0.1) \rangle, \\
 &\langle \check{B}_2, (0.32, 0.38), (0.1, 0.04) \rangle, \\
 &\langle \check{B}_3, (0.1, 0.36), (0.34, 0.12) \rangle, \langle \check{B}_4, (0.26, 0.15), (0.12, 0.2) \rangle \}.
 \end{aligned}$$

Then

$$\begin{aligned}
 \left((\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) \right)^c &= \{ \langle \check{B}_1, (0.24, 0.1), (0.2, 0.3) \rangle, \\
 &\langle \check{B}_2, (0.1, 0.04), (0.32, 0.38) \rangle, \\
 &\langle \check{B}_3, (0.34, 0.12), (0.1, 0.36) \rangle, \langle \check{B}_4, (0.12, 0.2), (0.26, 0.15) \rangle \}.
 \end{aligned}$$

Now consider R.H.S.

$$\begin{aligned}
 (\check{\delta}, \check{G}_1)^c \cap_\epsilon (\check{\delta}, \check{G}_2)^c &= \{ \langle \check{B}_1, (0.24, 0.1), (0.2, 0.3) \rangle, \\
 &\langle \check{B}_2, (0.1, 0.04), (0.32, 0.38) \rangle, \\
 &\langle \check{B}_3, (0.34, 0.12), (0.1, 0.36) \rangle, \langle \check{B}_4, (0.12, 0.2), (0.26, 0.15) \rangle \}.
 \end{aligned}$$

From this, it is clear that L.H.S.=R.H.S. Similarly, we can prove (2).

5.5 | Commutative Law

$$\begin{aligned}
 (\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) &= (\check{\delta}, \check{G}_2) \cup (\check{\delta}, \check{G}_1). \\
 (\check{\delta}, \check{G}_1) \cup_R (\check{\delta}, \check{G}_2) &= (\check{\delta}, \check{G}_2) \cup_R (\check{\delta}, \check{G}_1). \\
 (\check{\delta}, \check{G}_1) \cap (\check{\delta}, \check{G}_2) &= (\check{\delta}, \check{G}_2) \cap (\check{\delta}, \check{G}_1).
 \end{aligned}$$

$$(\check{\delta}, \check{G}_1) \cap_{\epsilon} (\check{\delta}, \check{G}_2) = (\check{\delta}, \check{G}_2) \cap_{\epsilon} (\check{\delta}, \check{G}_1).$$

Example 18. To *Prove (1)* law, we consider illustrative example. For this, suppose that

L.H.S:

$$\begin{aligned} (\check{\delta}, \check{G}_1) \cup (\check{\delta}, \check{G}_2) = \{ < \check{B}_1, (0.2, 0.3), (0.24, 0.1) >, \\ < \check{B}_2, (0.32, 0.38), (0.1, 0.04) >, \\ < \check{B}_3, (0.1, 0.36), (0.34, 0.12) >, < \check{B}_4, (0.26, 0.15), (0.12, 0.2) > \}. \end{aligned}$$

R.H.S:

$$\begin{aligned} (\check{\delta}, \check{G}_2) \cup (\check{\delta}, \check{G}_1) = \{ < \check{B}_1, (0.2, 0.3), (0.24, 0.1) >, \\ < \check{B}_2, (0.32, 0.38), (0.1, 0.04) >, \\ < \check{B}_3, (0.1, 0.36), (0.34, 0.12) >, < \check{B}_4, (0.26, 0.15), (0.12, 0.2) > \}. \end{aligned}$$

From above equation, we meet the required result. Similarly, we can prove the remaining.

6 | Conclusion

In this article, the basic fundamentals of refined intuitionistic fuzzy Set (RIFS) i.e. RIF subset, Equal RIFS, Complement of RIFS, Null RIFS and aggregation operators i.e. union, intersection, restricted intersection, extended union, extended intersection and restricted difference of two RIFS is defined. All these fundamentals are explained using an illustrative example. Further extension can be sought through developing similarity measures for comparison purposes.

References

- [1] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8, 338-353.
- [2] Zadeh, L. A. (1999). Fuzzy sets as a basis for a theory of possibility. *Fuzzy sets and systems*, 100, 9-34.
- [3] Dubois, D., & Prade, H. (1986). Fuzzy sets and statistical data. *European journal of operational research*, 25(3), 345-356.
- [4] Dubois, D., & Prade, H. (1991). Random sets and fuzzy interval analysis. *Fuzzy sets and systems*, 42(1), 87-101.
- [5] Dubois, D., & Prade, H. (1983). Ranking fuzzy numbers in the setting of possibility theory. *Information sciences*, 30(3), 183-224.
- [6] Beg, I., & Ashraf, S. (2009). Similarity measures for fuzzy sets. *Appl. comput. math*, 8(2), 192-202.
- [7] Beg, I., & Ashraf, S. (2008). Fuzzy similarity and measure of similarity with Lukasiewicz implicator. *New mathematics and natural computation*, 4(02), 191-206.
- [8] Beg, I., & Ashraf, S. (2009). Fuzzy inclusion and fuzzy similarity with gödel fuzzy implicator. *New mathematics and natural computation*, 5(03), 617-633.
- [9] Neog, T. J., & Sut, D. K. (2011). Complement of an extended fuzzy set. *International journal of computer applications*, 29(3), 39-45.
- [10] Uygun, O., Yalcin, S., Kiraz, A., & Furkan Erkan, E. (2020). A novel assessment approach to EFQM driven institutionalization using integrated fuzzy multi-criteria decision-making methods. *Scientia ironica*, 27(2), 880-892.
- [11] Damirchi-Darasi, S. R., Zarandi, M. F., Turksen, I. B., & Izadi, M. (2019). Type-2 fuzzy rule-based expert system for diagnosis of spinal cord disorders. *Scientia Iranica. Transaction E, industrial engineering*, 26(1), 455-471.

- [12] Brown, J. G. (1971). A note on fuzzy sets. *Information and control*, 18(1), 32-39.
- [13] Dubois, D., & Prade, H. (1989). Fuzzy sets, probability and measurement. *European journal of operational research*, 40(2), 135-154.
- [14] Dubois, D., & Prade, H. (1997). The three semantics of fuzzy sets. *Fuzzy sets and systems*, 90(2), 141-150.
- [15] Isah, A. I. (2020). Some algebraic structures of multi-fuzzy set. *Science world journal*, 15(1), 21-25.
- [16] Meng, F. Y., Tang, J., & Fujita, H. (2019). Consistency-based algorithms for decision-making with interval fuzzy preference relations. *IEEE transactions on fuzzy systems*, 27(10), 2052-2066.
- [17] Atanassov, K. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and systems*, 20 (1), 87-96.
- [18] Atanassov, K. T. (1999). Intuitionistic fuzzy sets. In *Intuitionistic fuzzy sets* (pp. 1-137). Physica, Heidelberg.
- [19] Ejegwa, P. A., Akowe, S. O., Otene, P. M., & Ikyule, J. M. (2014). An overview on intuitionistic fuzzy sets. *International journal of scientific and technology research*, 3(3), 142-145.
- [20] Szmidt, E., & Kacprzyk, J. (2001). Entropy for intuitionistic fuzzy sets. *Fuzzy sets and systems*, 118(3), 467-477.
- [21] Szmidt, E., & Kacprzyk, J. (2000). Distances between intuitionistic fuzzy sets. *Fuzzy sets and systems*, 114(3), 505-518.
- [22] Ersoy, B. A., & Davvaz, B. (2013, September). Structure of intuitionistic fuzzy sets in-semihyperring. In *Abstract and applied analysis* (Vol. 2013). Hindawi.
- [23] Bustince, H., & Burillo, P. (1996). Vague sets are intuitionistic fuzzy sets. *Fuzzy sets and systems*, 79(3), 403-405.
- [24] Garg, H., & Kaur, G. (2020). Extended TOPSIS method for multi-criteria group decision-making problems under cubic intuitionistic fuzzy environment. *Scientia iranica*, 27(1), 396-410.
- [25] Garg, H. (2016). A novel correlation coefficients between pythagorean fuzzy sets and its applications to decision-making processes. *International journal of intelligent systems*, 31(12), 1234-1252.
- [26] Ejegwa, P. A., & Onyeke, I. C. (2020). Medical diagnostic analysis on some selected patients based on modified Thao et al.'s correlation coefficient of intuitionistic fuzzy sets via an algorithmic approach. *Journal of fuzzy extension and applications*, 1(2), 130-141.
- [27] Edalatpanah, S. A. (2019). A data envelopment analysis model with triangular intuitionistic fuzzy numbers. *International journal of data envelopment analysis*, 7(4), 47-58.
- [28] Edalatpanah, S. A. (2020). Neutrosophic structured element. *Expert systems*. <https://doi.org/10.1111/exsy.12542>
- [29] Kumar, R., Edalatpanah, S. A., Jha, S., Gayen, S., & Singh, R. (2019). Shortest path problems using fuzzy weighted arc length. *International journal of innovative technology and exploring engineering*, 8(6), 724-731.
- [30] Gulzar, M., Alghazzawi, D., Mateen, M. H., & Kausar, N. (2020). A certain class of t-intuitionistic fuzzy subgroups. *IEEE access*, 8, 163260-163268.
- [31] Zhang, L., Zhan, J., & Yao, Y. (2020). Intuitionistic fuzzy TOPSIS method based on CVPIFRS models: an application to biomedical problems. *Information sciences*, 517, 315-339.
- [32] Mohan, S., Kannusamy, A. P., & Samiappan, V. (2020). A new approach for ranking of intuitionistic fuzzy numbers. *Journal of fuzzy extension and applications*, 1(1), 15-26.
- [33] Imeni, M. (2020). Fuzzy logic in accounting and auditing. *Journal of fuzzy extension and applications*, 1(1), 69-75.
- [34] Smarandache, F. (2019). Neutrosophic set is a generalization of intuitionistic fuzzy set, inconsistent intuitionistic fuzzy set (picture fuzzy set, ternary fuzzy set), pythagorean fuzzy set (atanassov's intuitionistic fuzzy set of second type), q-rung orthopair fuzzy set, spherical fuzzy set, and n-hyperspherical fuzzy set, while neutrosophication is a generalization of regret theory, grey system theory, and three-ways decision (revisited). Infinite Study.